

POLYCHLORINATED BIPHENYLS (Group 2A)

For definition of Groups, see [Preamble Evaluation](#).

Supplement 7: (1987) (p. 322)

CAS No.: 1336-36-3

Chem. Abstr. Name: Polychlorinated Biphenyls

Biphenyl:

CAS No.: 92-52-4

Chem. Abstr. Name: 1,1'-Biphenyl

2-Chlorobiphenyl:

CAS No.: 2051-60-7

Chem. Abstr. Name: 2-Chloro-1,1'-biphenyl

4-Chlorobiphenyl:

CAS No.: 2051-62-9

Chem. Abstr. Name: 4-Chloro-1,1'-biphenyl

2,2'-Dichlorobiphenyl:

CAS No.: 13029-08-8

Chem. Abstr. Name: 2,2'-Dichloro-1,1'-biphenyl

2,3'-Dichlorobiphenyl:

CAS No.: 25569-80-6

Chem. Abstr. Name: 2,3'-Dichloro-1,1'-biphenyl

2,4'-Dichlorobiphenyl:

CAS No.: 34883-43-7

Chem. Abstr. Name: 2,4'-Dichloro-1,1'-biphenyl

4,4'-Dichlorobiphenyl:

CAS No.: 2050-68-2

Chem. Abstr. Name: 4,4'-Dichloro-1,1'-biphenyl

2,2',3-Trichlorobiphenyl:

CAS No.: 38444-78-9

Chem. Abstr. Name: 2,2',3-Trichloro-1,1'-biphenyl

2,2',5-Trichlorobiphenyl:

CAS No.: 37680-65-2

Chem. Abstr. Name: 2,2',5-Trichloro-1,1'-biphenyl

2,3',4-Trichlorobiphenyl:

CAS No.: none available

Chem. Abstr. Name: 2,3',4-Trichloro-1,1'-biphenyl

2',3,4-Trichlorobiphenyl:

CAS No.: 38444-86-9

Chem. Abstr. Name: 2',3,4-Trichloro-1,1'-biphenyl

2,4,4'-Trichlorobiphenyl:

CAS No.: 7012-37-5

Chem. Abstr. Name: 2,4,4'-Trichloro-1,1'-biphenyl

2,4',5-Trichlorobiphenyl:

CAS No.: 16606-02-3

Chem. Abstr. Name: 2,4',5-Trichloro-1,1'-biphenyl

2,2'3,5'-Tetrachlorobiphenyl:

CAS No.: 41464-39-5

Chem. Abstr. Name: 2,2',3,5'-Tetrachloro-1,1'-biphenyl

2,2',4,5'-Tetrachlorobiphenyl:

CAS No.: 41464-40-8

Chem. Abstr. Name: 2,2',4,5'-Tetrachloro-1,1'-biphenyl

2,2',5,5'-Tetrachlorobiphenyl:

CAS No.: 35693-99-3

Chem. Abstr. Name: 2,2',5,5'-Tetrachloro-1,1'-biphenyl

2,3,4,4'-Tetrachlorobiphenyl:

CAS No.: 33025-41-1

Chem. Abstr. Name: 2,3,4,4'-Tetrachloro-1,1'-biphenyl

2,3',4,4'-Tetrachlorobiphenyl:

CAS No.: 32598-10-0

Chem. Abstr. Name: 2,3',4,4'-Tetrachloro-1,1'-biphenyl

2,3',4',5-Tetrachlorobiphenyl:

CAS No.: 32598-11-1

Chem. Abstr. Name: 2,3',4',5-Tetrachloro-1,1'-biphenyl

3,3',4,4'-Tetrachlorobiphenyl:

CAS No.: 32598-13-3

Chem. Abstr. Name: 3,3',4,4'-Tetrachloro-1,1'-biphenyl

2,2',3,3',6-Pentachlorobiphenyl:

CAS No.: 52663-60-2

Chem. Abstr. Name: 2,2',3,3',6-Pentachloro-1,1'-biphenyl

2,2',3,4,5'-Pentachlorobiphenyl:

CAS No.: 38380-02-8

Chem. Abstr. Name: 2,2',3,4,5'-Pentachloro-1,1'-biphenyl

2,2',3',4,5-Pentachlorobiphenyl:

CAS No.: 41464-51-1

Chem. Abstr. Name: 2,2',3',4,5-Pentachloro-1,1'-biphenyl

2,2',3,4',6-Pentachlorobiphenyl:

CAS No.: none available

Chem. Abstr. Name: 2,2',3,4',6-Pentachloro-1,1'-biphenyl

2,2',3,5',6-Pentachlorobiphenyl:

CAS No.: 38379-99-6

Chem. Abstr. Name: 2,2',3,5',6-Pentachloro-1,1'-biphenyl

2,2',4,4',5-Pentachlorobiphenyl:

CAS No.: 38380-01-7

Chem. Abstr. Name: 2,2',4,4',5-Pentachloro-1,1'-biphenyl

2,2',4,5,5'Pentachlorobiphenyl:

CAS No.: 37680-73-2

Chem. Abstr. Name: 2,2',4,5,5'-Pentachloro-1,1'-biphenyl

2,3,3',4,4'-Pentachlorobiphenyl:

CAS No.: 32598-14-4

Chem. Abstr. Name: 2,3,3',4,4'-Pentachlorol-1,1'-biphenyl

2,3,3',4',6-Pentachlorobiphenyl:

CAS No.: 38380-03-9

Chem. Abstr. Name: 2,3,3',4',6-Pentachlorol-1,1'-biphenyl

2,3',4,4',5-Pentachlorobiphenyl:

CAS No.: 31508-00-6

Chem. Abstr. Name: 2,3',4,4',5-Pentachlorol-1,1'-biphenyl

2,2',3,3',4,6-Hexachlorobiphenyl:

CAS No.: 38380-05-1

Chem. Abstr. Name: 2,2',3,3',4,6-Hexachloro-1,1'-biphenyl

2,2',3,3',6,6'-Hexachlorobiphenyl:

CAS No.: 38411-22-2

Chem. Abstr. Name: 2,2',3,3',6,6'-Hexachloro-1,1'-biphenyl

2,2',3,4,4',5-Hexachlorobiphenyl:

CAS No.: 35694-06-5

Chem. Abstr. Name: 2,2',3,4,4',5-Hexachloro-1,1'-biphenyl

2,2',3,4,4',5'-Hexachlorobiphenyl:

CAS No.: 35065-28-2

Chem. Abstr. Name: 2,2',3,4,4',5'-Hexachloro-1,1'-biphenyl

2,2',3',4,5,6'-Hexachlorobiphenyl:

CAS No.: 38380-04-0

Chem. Abstr. Name: 2,2',3',4,5,6'-Hexachloro-1,1'-biphenyl

2,2',4,4',5,5'-Hexachlorobiphenyl:

CAS No.: 35065-27-1

Chem. Abstr. Name: 2,2',4,4',5,5'-Hexachloro-1,1'-biphenyl

2,2',3,3',4,4',5-Heptachlorobiphenyl:

CAS No.: 35065-30-6

Chem. Abstr. Name: 2,2',3,3',4,4',5-Heptachloro-1,1'-biphenyl

2,2',3,3',4,5,6'-Heptachlorobiphenyl:

CAS No.: 38441-25-5

Chem. Abstr. Name: 2,2',3,3',4,5,6'-Heptachloro-1,1'-biphenyl

2,2',3,4,4',5,5'-Heptachlorobiphenyl:

CAS No.: 35065-29-3

Chem. Abstr. Name: 2,2',3,4,4',5,5'-Heptachloro-1,1'-biphenyl

A. Evidence for carcinogenicity to humans (*limited*)

Information on the possible carcinogenic risk of human exposure to polychlorinated biphenyls (PCBs) comes from studies of occupational populations and of populations exposed to the compounds accidentally. PCB mixtures may be contaminated with polychlorinated dibenzofurans and polychlorinated dibenzodioxins.

A slight increase in the incidence of cancer, particularly melanoma of the skin, was reported in a small group of men exposed to Aroclor 1254, a mixture of PCBs [ref: 1]. In a study of over 2500 US workers exposed to a similar mixture of PCBs during the manufacture of electrical capacitors, five deaths due to cancer of the liver and biliary passages were observed, whereas 1.9 would have been expected. This increase was sustained mainly by female workers in one of the two plants in the study (four of the five deaths), and all five had first been employed before the early 1950s [ref: 2,3]. Another study of workers in a capacitor plant was conducted in Italy. Exposure in the early years of production (until 1964) was to PCB mixtures containing 54% chlorine (mainly Aroclor 1254 and Pyralene 1476), which were later replaced by mixtures containing 42% chlorine (mainly Pyralene 3010 and 3011). Early results showed a significant excess of all cancers among male workers, which was due mainly to cancers of the digestive system and of the lymphatic and haematopoietic tissues. Among female workers, a slight increase in mortality from cancer of the lymphatic and haematopoietic tissues was reported [ref: 4]. The study was later enlarged and extended to include 2100 workers and to cover the period 1946-1982. Both male and female workers exhibited significantly increased cancer mortality in comparison with rates for the local population (14 observed, 7.6 expected; and 12 and 5.3, respectively, for men and women). Among male workers, cancers of the gastrointestinal tract (two stomach, two pancreas, one liver and one biliary passages) taken together were significantly increased (6 observed, 2.2 expected). Female workers showed a significant increase in deaths from haematological neoplasms (4 observed, 1.1 expected) [ref: 5]. In Sweden, among 142 male workers employed between 1965 and 1978 in a capacitor manufacturing plant when PCB mixtures containing up to 42% chlorine had been used, no significant excess of cancer deaths was noted. Cancer incidence was also examined: the number of cases observed corresponded well to that expected. One individual in a subgroup with higher exposure developed two relatively rare tumours, both of which occurred ten years after the start of exposure: a slow-growing mesenchymal tumour (desmoid) and a malignant lymphoma [ref: 6].

After contamination of cooking oil with a mixture of PCBs (Kanechlor 400) in Japan in 1968, a large population was intoxicated ('Yusho' disease). An early report on mortality from 1963-1983 showed a significantly increased risk of all cancers, and an almost five-fold significantly elevated risk of primary liver cancer. The edible rice oil had also been contaminated by polychlorinated quaterphenyls and polychlorinated dibenzofurans. Dose-response relationships were not clarified [ref: 7]. A further comprehensive study of 887 male 'Yusho' patients showed statistically significantly increased mortality from all malignancies (33 observed, 15.5 expected), from liver cancer (9 observed, 1.6 expected) and from lung cancer (8 observed, 2.5 expected). Use of local rather than national rates in calculating expected number of deaths decreased the observed:expected ratio for liver cancer from 5.6 to 3.9, which was still statistically significant. A closer look at the geographical distribution of liver cancer cases did not allow exclusion of factors other than PCB poisoning as a possible explanation for this finding. For the 874 female patients examined, none of the noted observed:expected ratios was significant [ref: 8]. In a series of ten autopsies of 'Yusho' patients, two adenocarcinomas of the liver were found, with no indication of a direct association with exposure to PCBs [ref: 9]. Ultrasonic and tumour marker examination of two series of 79 and 125 patients with 'Yusho' disease in 1983 and 1984, respectively, did not reveal any case of hepatic-cell carcinoma [ref: 10]. Two studies of the PCB content of fat tissues and cancer occurrence were available. An association was suggested between PCB concentrations in subcutaneous abdominal adipose tissue and the occurrence of cancers of the stomach, colon, pancreas, ovaries and prostate [ref: 11]. No indication emerged of a relationship between PCB content in extractable breast fat tissue and the occurrence of breast cancer [ref: 12].

The available studies suggest an association between cancer and exposure to PCBs. The increased risk from hepatobiliary cancer emerged consistently in different studies. Since, however, the numbers were small, dose-response relationships could not be evaluated, and the role of compounds other than PCBs could not be excluded, the evidence was considered to be limited.

B. Evidence for carcinogenicity to animals (*sufficient*)

Certain PCBs (particularly with greater than 50% chlorination) produced benign and malignant liver neoplasms in mice and rats after their oral administration [ref: 1,13,14]. Oral administration of Aroclor 1254 to rats yielded hepatocellular adenomas and carcinomas as well as intestinal metaplasia and a low, not statistically significant incidence of stomach adenocarcinomas [ref: 15]. PCBs were inadequately tested in mice for induction of skin tumours [ref: 16,17]. In several studies, oral or intraperitoneal administration of PCBs enhanced the incidences of preneoplastic lesions [ref: 18-20] and of neoplasms [ref: 21,22] of the liver induced in rats by *N*-nitrosodiethylamine or 2-acetylaminofluorene. In one study, intragastric administration of PCBs to mice increased the incidence of lung tumours induced by intraperitoneal administration of *N*-nitrosodimethylamine [ref: 23].

C. Other relevant data

No data were available on the genetic and related effects of PCBs in humans.

Dominant lethal effects were not induced in rats administered PCBs orally, but were produced in rats nursed by females that had received PCBs orally. PCBs did not induce chromosomal aberrations in bone-marrow cells or spermatogonia of rats treated *in vivo*; micronuclei were not induced in bone-marrow cells of mice in one study, while equivocal results were obtained in a second study in which the PCBs were administered in corn oil. They did not transform Syrian hamster embryo cells *in vitro*. PCBs induced DNA strand breaks and unscheduled DNA synthesis in rat hepatocytes *in vitro*.

Neither chromosomal breakage nor aneuploidy was induced in *Drosophila*. PCB mixtures did not induce SOS repair and were not mutagenic to bacteria [ref: 24].

2,2',5,5'-Tetrachlorophenyl induced DNA strand breaks in mouse cells *in vitro*.

2,4,5,2',4',5'-Hexachlorobiphenyl but not 3,4,5,3',4',5'-hexachlorobiphenyl inhibited intercellular communication in Chinese hamster V79 cells. Purified 2,4,2',4'-, 2,5,2',5'- and 3,4,3',4'-tetrachloro- and 2,4,6,2',4',6'-hexachlorobiphenyl were not mutagenic to bacteria [ref: 24].

Overall evaluation

Polychlorinated biphenyls are *probably carcinogenic to humans (Group 2A)*.

For definition of the italicized terms, see [Preamble Evaluation](#).

Also see previous evaluations: [Vol. 7 \(1974\)](#); [Vol. 18 \(1978\)](#)

References

1. IARC Monographs, 18, 43-103, 1978
2. Brown, D.P. & Jones, M. (1981) Mortality and industrial hygiene study of workers exposed to polychlorinated biphenyls. Arch. environ. Health, 36, 120-129
3. Brown, D.P. (1987) Mortality of workers exposed to polychlorinated biphenyls. An update. Arch. environ. Health (in press)
4. Bertazzi, P.A., Zocchetti, C., Guercilena, S., Della Foglia, M., Pesatori, A. & Riboldi, L. (1982) Mortality study of male and female workers exposed to PCB's. In: Proceedings of the International Symposium on Prevention of Occupational Cancer, Helsinki, 1981, Geneva, International Labour Office, pp. 242-248
5. Bertazzi, P.A., Riboldi, L., Pesatori, A., Radice, L. & Zocchetti, C. (1987) Cancer mortality of capacitor manufacturing workers. Am. J. ind. Med., 11, 165-176
6. Gustavsson, P., Hogstedt, C. & Rappe, C. (1986) Short-term mortality and cancer incidence in capacitor manufacturing workers exposed to polychlorinated biphenyls (PCBs). Am. J. ind. Med., 10, 341-344
7. Umeda, G. (1984) Studies on long-term effects of PCBs on human body: mortality from primary liver cancer and prevalence of ischemic heart disease (Abstract No. 19.8). In: Eustace, I.E., ed., Proceedings of the XXI International Congress on Occupational Health, Dublin, 1984, p. 166
8. Kuratsune, M., Nakamura, Y., Ikeda, M. & Hirohata, T. (1986) Analysis of deaths seen among patients with Yusho (Abstract FL17). In: Dioxin 86. Proceedings of the VI International Symposium on Chlorinated Dioxins and Related Compounds, Fukuoka, Japan, 1986, p. 179
9. Kikuchi, M. (1984) Autopsy of patients with Yusho. Am. J. ind. Med., 5, 19-30
10. Okumura, M. & Sakaguchi, S. (1985) Hepatic cell carcinoma and the patients with Yusho (Jpn.). Fukuoka Igaku Zasshi, 76, 229-232
11. Unger, M., Olsen, J. & Clausen, J. (1982) Organochlorine compounds in the adipose tissue of

- deceased persons with and without cancer: a statistical survey of some potential confounders. *Environ. Res.*, 29, 371-376
12. Unger, M., Kiaer, H., Blichert-Toft, M., Olsen, J. & Clausen, J. (1984) Organochlorine compounds in human breast fat from deceased persons with and without breast cancer and in a biopsy material from newly diagnosed patients undergoing breast surgery. *Environ. Res.*, 34, 24-28
13. Norback, D.H. & Weltman, R.H. (1985) Polychlorinated biphenyl induction of hepatocellular carcinoma in the Sprague-Dawley rat. *Environ. Health Perspect.*, 60, 97-105
14. Schaeffer, E., Greim, H. & Goessner, W. (1984) Pathology of chronic polychlorinated biphenyl (PCB) feeding in rats. *Toxicol. appl. Pharmacol.*, 75, 278-288
15. Ward, J.M. (1985) Proliferative lesions of the glandular stomach and liver in F344 rats fed diets containing Aroclor 1254. *Environ. Health Perspect.*, 40, 89-95
16. DiGiovanni, J., Viaje, A., Berry, D.L., Slaga, T.J. & Juchau, M.R. (1977) Tumor-initiating ability of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) and Aroclor 1254 in the two-stage system of mouse skin carcinogenesis. *Bull. environ. Contam. Toxicol.*, 18, 552-557
17. Hori, M., Fujita, K., Yamashiro, K., Toriyama, F., Hirose, R., Shukuwa, T., Toyoshima, H. & Yoshida, H. (1985) Methylcholanthrene induced mouse skin cancer. *Fukuoka Igaku Zasshi*, 76, 208-214
18. Pereira, M.A., Herren, S.L., Britt, A.L. & Khoury, M.M. (1982) Promotion by polychlorinated biphenyls of enzyme-altered foci in rat liver. *Cancer Lett.*, 15, 185-190
19. Buchmann, A., Kunz, W., Wolf, C.R., Oesch, F. & Robertson, L.W. (1986) Polychlorinated biphenyls, classified as either phenobarbital or 3-methylcholanthrene-type inducers of cytochrome P-450, are both hepatic tumor promoters in diethylnitrosamine-initiated rats. *Cancer Lett.*, 32, 243-253
20. Tatematsu, M., Nakanishi, K., Murasaki, G., Miyata, Y., Hirose, M. & Ito, N. (1979) Enhancing effect of inducers of liver microsomal enzymes on induction of hyperplastic liver nodules by N-2-fluorenylacetamine in rats. *J. natl Cancer Inst.*, 63, 1411-1416
21. Nishizumi, M. (1979) Effect of phenobarbital, dichlorodiphenyltrichloroethane, and polychlorinated biphenyls on diethylnitrosamine-induced hepatocarcinogenesis. *Gann*, 70, 835-837
22. Preston, B.D., Van Miller, J.P., Moore, R.W. & Allen, J.R. (1981) Promoting effects of polychlorinated biphenyls (Aroclor 1254) and polychlorinated dibenzofuran-free Aroclor 1254 on diethylnitrosamine-induced tumorigenesis in the rat. *J. natl Cancer Inst.*, 66, 509-515
23. Anderson, L.M., Ward, J.M., Fox, S.D., Isaaq, H.J. & Riggs, C.W. (1986) Effects of a single dose of polychlorinated biphenyls to infant mice on *N*-nitrosodimethylamine-initiated lung and liver tumors. *Int. J. Cancer*, 38, 109-116
24. IARC Monographs, Suppl. 6, 469-471, 1987

Synonyms for Polychlorinated biphenyls:

- Chlorinated biphenyl

Polychlorinated Biphenyls

- Chlorinated diphenyl
- Chlorobiphenyl
- PCB
- PCBs
- Polychlorinated biphenyl
- Polychlorobiphenyl

Trade names for Polychlorinated biphenyls (The producing country is given in parentheses):

- Aroclor (USA)
- Chlorextol (USA)
- Clophen (FRG)
- Dykanol (USA)
- Fenclor (Italy)
- Inerteen (USA)
- Kanechlor (Japan)
- Noflamol (USA)
- Phenoclor (France)
- Pyralene (as formulated, no longer contains PCBs) (France)
- Pyranol (USA)
- Santotherm (Japan)
- Sovol (USSR)
- Therminol (USA)

Last updated: 11 February 1998